

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.

1. REPORT DATE (DD-MM-YYYY)

2. REPORT TYPE
Technical Papers

3. DATES COVERED (From - To)

4. TITLE AND SUBTITLE

5a. CONTRACT NUMBER

5b. GRANT NUMBER

5c. PROGRAM ELEMENT NUMBER

6. AUTHOR(S)

5d. PROJECT NUMBER

2303

5e. TASK NUMBER

m2c8

5f. WORK UNIT NUMBER

7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)

Air Force Research Laboratory (AFMC)
AFRL/PRS
5 Pollux Drive
Edwards AFB CA 93524-7048

8. PERFORMING ORGANIZATION
REPORT

9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)

Air Force Research Laboratory (AFMC)
AFRL/PRS
5 Pollux Drive
Edwards AFB CA 93524-7048

10. SPONSOR/MONITOR'S
ACRONYM(S)

11. SPONSOR/MONITOR'S
NUMBER(S)

12. DISTRIBUTION / AVAILABILITY STATEMENT

Approved for public release; distribution unlimited.

13. SUPPLEMENTARY NOTES

14. ABSTRACT

15. SUBJECT TERMS

16. SECURITY CLASSIFICATION OF:

a. REPORT

Unclassified

b. ABSTRACT

Unclassified

c. THIS PAGE

Unclassified

17. LIMITATION
OF ABSTRACT

A

18. NUMBER
OF PAGES

19a. NAME OF RESPONSIBLE
PERSON

Leilani Richardson

19b. TELEPHONE NUMBER

(include area code)
(661) 275-5015

Standard Form 298 (Rev. 8-98)
Prescribed by ANSI Std. Z39.18

122 022

62

separate items are each

C8

MEMORANDUM FOR PRS (Contractor Publication)

FROM: PROI (TI) (STINFO)

25 October 2000

SUBJECT: Authorization for Release of Technical Information, Control Number: **AFRL-PR-ED-TP-2000-206**
Boatz, Jerry; Thompson, D. (OK State Univ), "Bond Dissociation Energies of Energetic Compounds: A Comparison of Theoretical Methods"

HEDM Contractors Conference (Park City, UT, 23-26 Oct 2000)
(Deadline: 20 Oct 2000 - PAST DUE)

(Statement A)

1. This request has been reviewed by the Foreign Disclosure Office for: a.) appropriateness of distribution statement, b.) military/national critical technology, c.) export controls or distribution restrictions, d.) appropriateness for release to a foreign nation, and e.) technical sensitivity and/or economic sensitivity.

Comments: _____

Signature _____ Date _____

2. This request has been reviewed by the Public Affairs Office for: a.) appropriateness for public release and/or b) possible higher headquarters review

Comments: _____

Signature _____ Date _____

3. This request has been reviewed by the STINFO for: a.) changes if approved as amended, b.) appropriateness of distribution statement, c.) military/national critical technology, d.) economic sensitivity, e.) parallel review completed if required, and f.) format and completion of meeting clearance form if required

Comments: _____

Signature _____ Date _____

4. This request has been reviewed by PRS for: a.) technical accuracy, b.) appropriateness for audience, c.) appropriateness of distribution statement, d.) technical sensitivity and economic sensitivity, e.) military/national critical technology, and f.) data rights and patentability

Comments: _____

APPROVED/APPROVED AS AMENDED/DISAPPROVED

PHILIP A. KESSEL
Technical Advisor

Date

Cleared (PA) _____
Logged (PA) _____
Notified (PA) _____
Copied & Distributed (STINFO) _____
This original is for PA files

Bond Dissociation Energies of Energetic Compounds: A Comparison of Theoretical Methods

Jerry A. Boatz

Air Force Research Laboratory
Propulsion Sciences and Advanced Concepts Division
AFRL/PRSP
Edwards AFB, CA 93524

and

Donald L. Thompson
Department of Chemistry
Oklahoma State University
Stillwater, OK 74078

20021122 022

AFOSR High Energy Density Matter Contractors' Conference
Park City, UT
24-26 Oct 00

**Bond Dissociation Energies of Energetic Compounds:
A Comparison of Theoretical Methods**

DISTRIBUTION STATEMENT A:
Approved for Public Release -
Distribution Unlimited

Jerry Boatz
Air Force Research Laboratory
AFRL/PRSP
10 E. Saturn Blvd.
Edwards AFB, CA 93524-7680

Donald L. Thompson
Department of Chemistry
Oklahoma State University
Stillwater, OK 74078

ABSTRACT

Due to their computational efficiency, density functional methods utilizing semiempirical hybrid functionals such as B3LYP are commonly used in calculating the molecular potential energy surfaces of a wide variety of molecules. In particular, potential energy surfaces of several energetic compounds such as hexahydro-1,3,5-trinitro-1,3,5-triazine (RDX), 1,3,3-trinitroazetidine (TNAZ), 5-nitro-2,4-dihydro-3*H*-1,2,4-triazolone (NTO), and 1,1-diamino-2,2-dinitroethylene recently have been computed using density functional methods (B3LYP). These potential energy surfaces are used, for example, to predict the decomposition mechanisms and for constructing force fields for modeling and simulation of solid state phase transitions.

The accuracy of B3LYP relative energies and structures is generally observed to be comparable to that obtained using second order perturbation theory methods. However, an unresolved issue is the reliability of density functional based methods in cases where a significant degree of multiconfigurational character may be present in the compounds of interest, such as in the energetic molecules listed above.

The present study is a systematic comparison of several electronic structure methods in the prediction of bond dissociation energies of energetic molecules. In particular, the C-NO₂ and C-NH₂ bond dissociation energies of 1,1-diamino-2,2-dinitroethylene and its prototypes are used as a testbed for comparison of B3LYP with single configuration self-consistent field (SCF), second order perturbation theory (MP2), and coupled-cluster (CCSD(T)) calculations as well as multiconfigurational SCF (MCSCF) and quasi-dependent perturbation theory (MCQDPT) methods.

GOAL

Increase propellant specific impulse by 10-20%.

OBJECTIVES

Discover, characterize, and develop new high energy density chemical propellants.

TECHNICAL CHALLENGES

Energetic compounds are frequently unstable.

Synthetic routes are unknown.

Limited chemical tuition to guide choice of targets.

APPROACH

Predict energy content and kinetic stability characteristics using computational chemistry.

Determine which theoretical method(s) are best suited for prediction of reliable energetics.

OVERVIEW

One of the most widely used theoretical methods for computing properties of energetic molecules is density functional theory, due to its computational efficiency and the general accuracy of the results (comparable to second order perturbation theory).

DFT is generally benchmarked against other QC methods by comparing predicted properties of selected molecules with a modest amount of multiconfigurational character.

ISSUES

How accurate are DFT predictions (e.g., bond energies) for molecules with a high degree of multiconfigurational character?

How does one determine if a molecule has a high degree of multiconfigurational character?

- Large spin contamination for open-shell species
- Negative LUMO energy for neutral closed shells.
- Natural orbital occupation numbers $> \sim 0.1$ for “virtual” orbitals.
- CCSD(T) “T1” diagnostic.
- Non-physical NOONs[1].

METHODOLOGY


Use MBPT, CCSD(T) non-physical (i.e., > 2 or < 0) natural orbital occupation numbers[1] (NOON) as diagnostic for multiconfigurational character.

Use predicted C-NH₂ and C-NO₂ bond dissociation energies of 1,1-diamino-2,2-dinitroethylene ("FOX-7") and its simpler prototypes as testbed.

Compare DFT with other single reference methods (MBPT2, CCSD(T), G2) and with multireference methods (MCSCF, MCQDPT).





1. M.S. Gordon, M.W. Schmidt, G.M. Chaban, K. R. Glaesemann, W.J. Stevens, and C. Gonzalez, "A Natural Orbital Diagnostic for Multiconfigurational Character in Correlated Wavefunctions", J. Chem. Phys., 110, 4199 (1999).

FOX-7: a “Push-Pull” Ethylene



Multiple Lewis valence structures suggest that FOX-7 may require multiconfigurational methods.

DFT Geometries

	C=C	C-NH ₂	C-NO ₂
	1.331* 1.327***		
	1.340* 1.336***	1.386* 1.392***	
	1.326* 1.322***		1.459* 1.473***
	1.426* 1.428** 1.421*** (1.456)	1.339* 1.345** 1.345*** (1.319, 1.325)	1.424* 1.432** 1.438*** (1.399, 1.426)


* P. Politzer, M.C. Concha, M.E. Grice, J.S. Murray, P. Lane, and D. Habibollahzadeh
Theochem, 452, 75(1998)

** A. Gindulyte, L. Massa, L. Huang, and J. Karl, J. Phys. Chem. A, 103, 11045(1999)

*** This work (B3LYP/6-311G(d,p))

() U. Benm and H. Östmark, Acta Cryst. C54, 1997(1998).

A Preliminary test: the CH₂ Bend potential




Circles: B3LYP/aug-cc-pVTZ


Curves: aug-cc-pVTZ basis used for all methods; taken from M.S. Gordon, M.W. Schmidt, G.M. Chaban, K. R. Glaesemann, W.J. Stevens, and C. Gonzalez, "A Natural Orbital Diagnostic for Multiconfigurational Character in Correlated Wavefunctions", J. Chem. Phys., 110, 4199 (1999).

Non-physical Natural Orbital Occupation Numbers


MBPT(2) CCSD(T)




-0.0001 (2)



-0.00014



2.296, 2.12, 2.00001
-0.99092, -0.xxxx




2.07575
-0.09409



2.00001

C-NO₂ Bond Dissociation Energies (kcal/mol)

	X=H	X=NH ₂	X=NH ₂	X=NH ₂
	Y=H	Y=H	Y=NH ₂	Y=NH ₂
	Z=H	Z=H	Z=H	Z=NO ₂
DFT	68.8	81.4		67.4

MBPT(2)

77.7


MCSCF

G2(MP2)

CCSD(T)//MBPT2

MCQDPT(2)//MCSCF

C-NH₂ Bond Dissociation Energies (kcal/mol)



DFT 105.2

MBPT(2) 115.6

MCSCF

G2(MP2)

CCSD(T)//MBPT2

MCQDPT(2)//MCSCF

SUMMARY

1. The bend potential of 1A_1 CH₂ has been computed using DFT (B3LYP) and compared against earlier MBPT(2), MCQDPT(2), CCSD(T), and MRCI calculations.
2. The B3LYP C-NO₂ and C-NH₂ bond dissociation energies of FOX-7 and its prototypes have been benchmarked against correlated single-configuration (MBPT(2), CCSD(T)) and multiconfigurational methods (MCSCF, MCQDPT).

CONCLUSIONS

1. The B3LYP 1A_1 CH₂ bend potential is in good agreement with MRCI, CCSD(T), and MCQDPT(2) predictions. It is in nearly exact agreement with MBPT(2).
2. B3LYP gives C-NH₂ bond dissociation energies in FOX-7 within ~5 kcal/mol of G2(MP2), CCSD(T), and MCQDPT(2).
3. B3LYP apparently “captures” a reasonable amount of non-dynamical correlation energy (e.g., works well for 2-configuration problems.)
4. B3LYP performance for C-NO₂ BDEs: TBD.

FUTURE DIRECTIONS

1. Examine effects of more flexible basis sets (e.g., cc-pvtz).
2. Benchmark DFT methods using other highly multiconfigurational molecules (e.g., FOOF).

ACKNOWLEDGEMENTS

Dr. Jeff Sheehy

Army Research Laboratory Major Shared
Resource Center